

What is a Degree Distribution?

Sofia C. Olhede and Patrick J. Wolfe*

November 29, 2012

Abstract

The most studied aspect of statistical network models is their degree structure, reflecting the propensity of nodes within a network to form connections with other nodes. Yet many simple random graph models are understood only asymptotically; for finitely many nodes, they do not yield either closed-form statistical likelihoods or precise forward generating mechanisms. In contrast, we provide exact statistical results, limit theorems, and large-sample approximations that govern the behavior of networks based on random weights whose pairwise products parameterize independent Bernoulli trials. For power-law degree sequences we make the important observation that the frequently observed exponential cutoff behavior can be explained as an effect of this sampling. This enables us for the first time to understand, from a statistical perspective, the heterogeneity of network degrees observed in practice, and to explore how properties of degree sequences scale with network size. Our results thus provide new tools to quantify degree variation within and across network populations.

Key words: Complex networks, power laws, random graphs

Network models play many important roles across the sciences, and have recently given rise to new tools for statistical inference [1, 2] in fields ranging from biology [3] to computational social science [4]. The single most studied aspect of network models is their degree structure, reflecting the propensity of nodes within a network to form connections with other nodes. At the same time, scaling is a fundamental property of physical processes, including those that give rise to networks [5]. As we collect larger and more complex network datasets, the scaling properties of network degree sequences must therefore be quantified and understood, to better enable the reproduction of characteristics seen in these data.

*The authors are with the Departments of Statistical Science and Computer Science, University College London. Work supported in part by the US Army Research Office under PECASE Award W911NF-09-1-0555 and by the US Office of Naval Research under MURI Award 58153-MA-MUR, and by the UK EPSRC under a Mathematical Sciences Leadership Fellowship and Institutional Sponsorship Award, and the UK Royal Society under a Wolfson Research Merit Award.

Thus we must develop a precise *statistical* understanding of the notion of a degree distribution. To do so we work with a hierarchical network model construction that encompasses many known random graph models [6, 7, 8], while at the same time allows for new types of heterogeneous structure and likelihood-based statistical inference. Network degree structure is regulated both by scaling and by the mechanism responsible for network generation. Starting with a model that is fully determined by its degree sequence, we parameterize the probabilities governing these degrees in order to obtain a forward generating mechanism. This fully stochastic model is useful and important because it enables a heterogeneous *population* of networks to be realized through a single, self-consistent framework.

1 Modeling the Degree Sequence of a Network

1.1 Existing Models for Network Degrees

The preeminent summary statistic of an observed network is its degree sequence. A natural modeling approach is to start by either fixing or sampling a length- n vector $\mathbf{d}$ of positive integer degrees [6], and then to select uniformly at random from the set of all possible graphs on n nodes that admit $\mathbf{d}$ as their degree sequence. In practice, edges are sequentially assigned to nodes in ways that respect $\mathbf{d}$ akin to sampling with replacement [9]. Asymptotically as $n \rightarrow \infty$ this practical construction is valid, but it can result in graphs with loops and multiple edges [10].

An important modification relaxes the requirement of a specified $\mathbf{d}$, replacing it with a set of n parameters describing the propensity of each node to form connections. Edges are then independent Bernoulli trials with success probabilities p_{ij} for all $1 \leq i \leq j \leq n$. Chung & Lu [11] associated a nonnegative weight w_i to each node i and set $p_{ij} = w_i w_j / \|\mathbf{w}\|_1$, where $\|\mathbf{w}\|_1 = \sum_{k=1}^n w_k$. Under the normalization constraint that $w_i^2 \leq \|\mathbf{w}\|_1$ for all i , it follows that the expected degree $\mathbb{E}(d_i)$ of the i th node is equal to w_i , and thus the unnormalized weights $w_1, \dots, w_n$ can be interpreted as expected degrees. This model is immediately recognizable as a log-linear model $\log p_{ij} = \log w_i + \log w_j - \log \|\mathbf{w}\|_1$, connecting to existing statistical methods such as maximum likelihood estimation of p_{ij} from an observed network [2, 12].

The given expected degrees model of [11] can be thought of as a special case of assigning edge probabilities p_{ij} using a kernel [8, Section 16.4]; see also [13, 14], where limiting properties of the resultant networks are derived. Further statistical insight was given by Chatterjee, Diaconis & Sly [15], who (among others [7, 8]) observed that under a similar model with $\log[p_{ij}/(1 - p_{ij})] = \tilde{w}_i + \tilde{w}_j$, the network degrees are a sufficient statistic for estimation of $\tilde{\mathbf{w}}$ —meaning that a network appears in the model’s statistical likelihood function only through its degrees. This model has the advantage of assigning equal likelihood to all graphs with the same degree sequence, though it loses the multiplicative structure of [11]. Janson [13] gives conditions under which these two models are

asymptotically equivalent as $n \rightarrow \infty$, and Perry & Wolfe [12] give graph sparsity conditions for fixed n under which optimal likelihood-based estimates for both models can be obtained directly as $\hat{p}_{ij} = d_i d_j / \|\mathbf{d}\|_1$ —a natural estimator based on Chung & Lu’s formulation [11], and for which we develop a limit theorem below. This degree-based estimator already sees wide use in practice, most predominantly in the calculation of modularity statistics for community detection in complex networks [1, 16].

1.2 Parameterizing the Poisson–Binomial Degree Distribution

Simple n -node graphs are represented by a symmetric adjacency matrix $\mathbf{A}$:

$$A_{ij} = \text{Bernoulli}(p_{ij}), 1 \leq i < j \leq n; A_{ji} = A_{ij}, A_{ii} = 0. \quad (1)$$

Each network degree is a sum $d_i = \sum_{j \neq i} A_{ij}$ of $n-1$ independent Bernoulli(p_{ij}) variates, with $p_{ji} = p_{ij}$, and thus takes the Poisson–Binomial distribution on values $k \in \{0, 1, \dots, n-1\}$:

$$\mathbb{P}(d_i = k \mid \{p_{ij}\}_{j \neq i}) = \sum_{l=1}^{\binom{n-1}{k}} \left(\prod_{j' \in A_i^l} p_{ij'} \right) \left(\prod_{j' \in \bar{A}_i^l} (1 - p_{ij'}) \right). \quad (2)$$

Here the $\binom{n-1}{k}$ sets A_i^l index, for fixed i , all distinct ways of obtaining k successes out of the $n-1$ Bernoulli trials $\{A_{ij}\}_{j \neq i}$, and each $\bar{A}_i^l$ is the respective complement in $\{0, 1, \dots, n-1\}$. Setting $p_{ij} = p$ for all $1 \leq i < j \leq n$ recovers the canonical random graph model studied by Erdős, Rényi, and others, in which case (2) reduces to the Binomial($n-1, p$) distribution for all i .

To fully understand degree distributions from a statistical point of view, we must be able to obtain a precise specification of degree behavior for any fixed network size n , as well as to develop large-sample properties of sequences of networks as n increases. For this reason we adopt the multiplicative model of [11], but with a parameterization that enables *individual* constraints to be specified for a *normalized* weight vector $\boldsymbol{\pi}$:

$$p_{ij} = \pi_i \pi_j, 1 \leq i < j \leq n; \pi_i \in \Pi_n \subseteq [0, 1], i = 1, \dots, n. \quad (3)$$

Note that relative to the parameterization $p_{ij} = w_i w_j / \|\mathbf{w}\|_1$, the formulation of (3) *decouples* the edge generation probabilities p_{ij} , so that each depends only on two parameters, and dispenses with the need to constrain $w_i^2 \leq \|\mathbf{w}\|_1$ for all i . This simple reformulation has important consequences for our statistical understanding of network models for fixed and growing n , and will later enable us to introduce a *hierarchical* model where the normalized weights $\pi_1, \dots, \pi_n$ are considered as a set of statistically *independent* random variables.

Thus far we have established that, conditional upon a weight vector $\boldsymbol{\pi}$, each network degree d_i follows a Poisson–Binomial distribution $\mathbb{P}(d_i = k \mid \boldsymbol{\pi})$ specified by (2). To understand the scaling properties of network degrees as $n \rightarrow \infty$, we

will allow the range Π_n of each normalized weight in (3) to depend on n . Fixing $\Pi_n = [0, 1]$ for all n ensures that elements of $\boldsymbol{\pi}$ can remain order one, while letting $|\Pi_n|$ shrink with n will reduce the order of each p_{ij} as $n \rightarrow \infty$.

To understand the effects of this n -dependent scaling, first observe that together (1)–(3) imply, via direct calculation, the following mean, variance, and covariance expressions for network degrees:

$$\mathbb{E}(d_i | \boldsymbol{\pi}) = \sum_{j \neq i} \mathbb{E}(A_{ij} | \boldsymbol{\pi}) = \pi_i \sum_{j \neq i} \pi_j, \quad (4)$$

$$\text{Var}(d_i | \boldsymbol{\pi}) = \sum_{j \neq i} \text{Var}(A_{ij} | \boldsymbol{\pi}) = \mathbb{E}(d_i | \boldsymbol{\pi}) - \pi_i^2 \sum_{j \neq i} \pi_j^2, \quad (5)$$

$$\text{Cov}(d_i, d_j | \boldsymbol{\pi}) = \text{Var}(A_{ij} | \boldsymbol{\pi}) = \pi_i \pi_j (1 - \pi_i \pi_j), \quad i \neq j. \quad (6)$$

The average behavior of degree d_i in n thus depends on its normalized weight π_i scaled by the sum $\|\boldsymbol{\pi}\|_1$, while its variance depends on the scaling $\|\boldsymbol{\pi}\|_1 - \pi_i \|\boldsymbol{\pi}\|_2^2$. The covariance between distinct degrees, on the other hand, cannot exceed $1/4$, and so the correlation between any two degrees in a single network realization will decay toward zero whenever $\text{Var}(d_i | \boldsymbol{\pi})$ is growing in n for each i . This result is well known for various random graph models in the literature; the interplay among (4)–(6) shows explicitly why and how rapidly (in n) this decorrelation occurs, as a function of our assumptions on the growth of $\|\boldsymbol{\pi}\|_1$ in n .

1.3 Degree Distribution Variability: Power Laws and Network Scaling

For any fixed n and weight vector $\boldsymbol{\pi}$, the methods above precisely quantify degree variability, in accordance with the Poisson–Binomial distribution $\mathbb{P}(d_i = k | \boldsymbol{\pi})$ of (2). The scaling of network behavior in n is equally critical to understanding the variability of the resulting network degrees. In general $\|\boldsymbol{\pi}\|_2^2$ and $\|\boldsymbol{\pi}\|_1$ will grow at different rates in n , and their ratio will impact not only the degree of heterogeneity incurred from Bernoulli sampling under the likelihood specified by (1) and the multiplicative model of (3), but also important aspects of statistical data analysis, such as our ability to estimate $\boldsymbol{\pi}$ from a single observed network under this combined model.

A natural example is given by graphs having power laws as limiting degree distributions, as illustrated in Fig. 1 below. Here the expected proportion of degree- k nodes in a single network realization scales as $k^{-\beta}$ for some power $\beta > 0$. Our first theorem treats power-law networks corresponding to those studied in [17]. It characterizes their degree sequences under the parameterization of (3), both when n is large and fixed, and asymptotically as $n \rightarrow \infty$.

To interpret this theorem, we need the following notation. Given two sequences f_n and g_n , the notation $f_n = \omega(g_n)$ means that f_n grows in n *strictly faster* than g_n , so that eventually (in n) the ratio $|g_n/f_n|$ becomes arbitrarily close to zero. In contrast, $f_n = \mathcal{O}(g_n)$ means that f_n grows *no faster* in n than g_n , in that $|f_n/g_n|$ is eventually upper bounded by a constant as $n \rightarrow \infty$.

Finally, $f_n = o(g_n)$ means that f_n grows *strictly slower* in n than g_n , so that $|f_n/g_n|$ eventually becomes arbitrarily close to zero.

Theorem 1 (Power law degrees). *Fix an exponent $\gamma \in (0, 1)$ and set*

$$\pi_i = \theta_n i^{-\gamma}, \quad i = 1, 2, \dots, n,$$

with θ_n an n -dependent sequence of scaling constants satisfying $\theta_n \leq 1$ for all n , so that $\Pi_n = [0, \theta_n]$ in the model of (3).

Then the expected value of the i th degree $d_i | \boldsymbol{\pi}$ is given by

$$\mathbb{E}(d_i | \boldsymbol{\pi}) = \theta_n^2 i^{-\gamma} [(1 - \gamma)^{-1} n^{1-\gamma} + \mathcal{O}(1)], \quad (7)$$

and as $n \rightarrow \infty$, each $d_i | \boldsymbol{\pi}$ approaches a Poisson random variable with mean $\mathbb{E}(d_i | \boldsymbol{\pi})$. In particular, the sum of all n differences

$$\left| \mathbb{P}(d_i = k | \boldsymbol{\pi}) - \frac{\mathbb{E}(d_i | \boldsymbol{\pi})^k e^{-\mathbb{E}(d_i | \boldsymbol{\pi})}}{k!} \right|, \quad k = 0, 1, \dots, n-1,$$

converges to zero, and thus so does each individual difference in turn.

Whenever $\mathbb{E}(d_i | \boldsymbol{\pi})$ is growing in n as determined by (7), then d_i becomes close to its expectation in that $d_i / \mathbb{E}(d_i | \boldsymbol{\pi})$ converges in probability to 1, and $(d_i - \mathbb{E}(d_i | \boldsymbol{\pi})) / \sqrt{\mathbb{E}(d_i | \boldsymbol{\pi})}$ converges in distribution to a Normal(0, 1) random variable.

Finally, assume $\gamma \in (0, 1/2)$ and $\theta_n = \omega(\sqrt{\log n} / n^{1/2-\gamma})$, in which case each degree d_i will grow in n . In expectation, an observed degree d_j from a single network realization, with j sampled uniformly from $\{1, \dots, n\}$, will obey for $\beta = 1 + 1/\gamma$ the power law

$$\mathbb{P}(d_j > x) = \left(\frac{x}{\mathbb{E}(d_n | \boldsymbol{\pi})} \right)^{1-\beta} \left[1 + \mathcal{O} \left(\frac{\sqrt{\log n}}{\sqrt{x}} \right) \right] + \mathcal{O} \left(\frac{1}{n^{2\alpha}} \right),$$

with $\alpha = 1 + o(1)$, for $x \in \{\mathbb{E}(d_n | \boldsymbol{\pi}), \mathbb{E}(d_{n-1} | \boldsymbol{\pi}), \dots, \mathbb{E}(d_2 | \boldsymbol{\pi})\}$.

Proof. Approximating $\|\boldsymbol{\pi}\|_1 = \theta_n n^{-\gamma} \sum_{i=1}^n (i/n)^{-\gamma}$ by a integral squeezing argument gives $\|\boldsymbol{\pi}\|_1 = \theta_n [n^{1-\gamma}/(1-\gamma) + \mathcal{O}(1)]$ for all $\gamma \in (0, 1)$; combining with (4) yields the expression for $\mathbb{E}(d_i | \boldsymbol{\pi})$.

In the same manner, we may verify that $\lim_{n \rightarrow \infty} [1 - \text{Var}(d_i | \boldsymbol{\pi}) / \mathbb{E}(d_i | \boldsymbol{\pi})] = 0$. For every $n = 1, 2, \dots$, this quantity controls the total variation distance between the Poisson–Binomial distribution of (2) and a Poisson distribution with mean $\mathbb{E}(d_i | \boldsymbol{\pi})$ [18], which is equivalent to the theorem statement.

Next, observe that if $\mathbb{E}(d_i | \boldsymbol{\pi})$ is growing in n , it follows from the limit above that $\lim_{n \rightarrow \infty} \text{Var}(d_i | \boldsymbol{\pi}) / \mathbb{E}(d_i | \boldsymbol{\pi})^2 = 0$, implying via Chebyshev's inequality that $d_i / \mathbb{E}(d_i | \boldsymbol{\pi})$ converges in probability to 1. Growth of $\mathbb{E}(d_i | \boldsymbol{\pi})$ implies growth of $\text{Var}(d_i | \boldsymbol{\pi})$ —a condition sufficient for the Lindeberg–Feller central limit theorem to apply.




Figure 1: Simulated degrees illustrating within- and between-network degree variability for a power-law network with exponent $\beta = 3$. The left-hand panel orders the degrees by number i , while the right-hand panel orders the degrees by size.

For the empirical distribution of n observed degrees, we write $F_n(x, \mathbf{d} | \boldsymbol{\pi}) = 1 - n^{-1} \sum_{i=1}^n \mathbb{I}(d_i > x | \boldsymbol{\pi})$, with the indicator function $\mathbb{I}(\cdot)$ taking values 0 or 1 accordingly. We may then compute, for x in the given range of the theorem, $\mathbb{E}(n^{-1} \sum_{i=1}^n \mathbb{I}(d_i > x | \boldsymbol{\pi})) = n^{-1} \sum_{i=1}^n \mathbb{P}(d_i > x | \boldsymbol{\pi})$ as follows. For those i such that $|x - \mathbb{E}(d_i | \boldsymbol{\pi})| \geq 2\sqrt{\mathbb{E}(d_i | \boldsymbol{\pi}) \ln n}$, the components $\mathbb{P}(d_i > x | \boldsymbol{\pi})$ in this sum can be treated using Chernoff bounds for tail probabilities of Poisson–Binomial variates. Contributions from the remaining values of i become increasingly negligible as n grows large, and thus can be bounded. Collecting the corresponding error terms together then yields the stated result for $n^{-1} \sum_{i=1}^n \mathbb{P}(d_i > x | \boldsymbol{\pi})$. $\square$

In addition to verifying power law behavior under the network adjacency model $A_{ij} = \text{Bernoulli}(\pi_i \pi_j)$, Theorem 1 reveals that the notion of a degree distribution has *two distinct interpretations* that can lead to radically different assessments of degree behavior. Figure 1 illustrates this by way of network realizations for the case $\beta = 3$, corresponding to the well-studied preferential attachment network model of [5]. The first result of Theorem 1 confirms the *between-network* variability of the i th degree as becoming Poissonian across an *ensemble* of network realizations; we can visualize this ensemble via the left-hand panel of Fig. 1, by looking across these realizations for any fixed i . Here the Bernoulli trials governing edge formation are the only source of variation. By contrast, the final result of Theorem 1 (which can be directly extended to the case $\beta = 3$) describes the expected *within-network* degree variability in terms of a power law. Here we see variability induced by the range of values of π_i , emphasized by the ordering of degrees within each realization as in the right-hand panel of Fig. 1.

In the special case of an Erdős–Rényi graph, the above interpretations agree. Here all π_i are equal rather than decreasing in i , and thus Theorem 1 as written does not apply. Uniquely in this special case, the degrees are identically distributed. Their correlation is furthermore equal to $1/(n-1)$, and thus as n becomes large, they decorrelate and behave like a random sample from $\mathbb{P}(d_i = k | \boldsymbol{\pi})$ —equivalently, like an ensemble of independent and identically distributed degree realizations. Later we will see that a similar construction can be used to produce a wide range of desired degree behavior, by letting elements of $\boldsymbol{\pi}$ themselves comprise a random sample.

More generally, as long as each $\text{Var}(d_i | \boldsymbol{\pi})$ is growing in n , then all degrees within a single network decorrelate in n , allowing us to view a single realized degree sequence as representing an ensemble of essentially independent—though *not* identically distributed—degree realizations. This view of a network’s set of realized degrees as a distribution prevails in the complex networks literature: it is a natural and meaningful empirical summary of *within-network* degree variability that respects the lack of any given ordering of the degrees.

In this view we consider a set of observed or expected degrees associated with a single network realization, and summarize their empirical distribution as a histogram. Note however that the observed degrees are random, while the expected degrees are not. Thus when reasoning about observed network data, we must take between-network sampling variability into account. This is why we could not simply consider the distribution function of the *expected* degrees, $F_n(x, \mathbb{E}(\mathbf{d} | \boldsymbol{\pi}))$, in the proof of Theorem 1 and invert F_n to solve $\mathbb{E}(d_i | \boldsymbol{\pi}) = [1 - F_n]^{-1}(i/n)$ in the limit of large n . Doing so verifies that when degree index j is sampled uniformly at random, $\mathbb{P}(\mathbb{E}(d_j | \boldsymbol{\pi}) > k) \propto k^{-\beta}$, whereas the arguments of Theorem 1 are required to ensure that the *observed degrees from a single network*, taken together as a group, will exhibit this same expected behavior.

1.4 A Power Law Central Limit Theorem

We now turn our attention to estimation of $\boldsymbol{\pi}$ from an observed network. Recall that in the power-law setting of Theorem 1, whenever an expected degree $\mathbb{E}(d_i | \boldsymbol{\pi})$ grows in n , the limiting Poisson distribution of d_i will tend toward a Normal distribution. As we show below, when n is large and the estimator $\hat{\pi} = d_i / \sqrt{\|\mathbf{d}\|_1}$ is used to estimate π_i [12], then $\hat{\pi}_i$ is approximately distributed as a $\text{Normal}(\pi_i, \pi_i / \|\boldsymbol{\pi}\|_1)$ random variable.

Theorem 2 (Power law central limit theorem for $\boldsymbol{\pi}$). *Assume as in Theorem 1 that each $\pi_i = \theta_n i^{-\gamma}$, with $\gamma \in (0, 1)$. Define $\hat{\pi}_i = d_i / \sqrt{\|\mathbf{d}\|_1}$ as an estimator of π_i , and assume $\mathbb{E}(d_i | \boldsymbol{\pi})$ is growing in n . Then as $n \rightarrow \infty$, the standardized variate $(\hat{\pi}_i - \pi_i) / \sqrt{\pi_i / \|\boldsymbol{\pi}\|_1}$ converges in distribution to a $\text{Normal}(0, 1)$ random variable.*

Proof. The growth of $\mathbb{E}(d_i | \boldsymbol{\pi})$ in n drives this result. Expand $(\hat{\pi}_i - \pi_i) / \sqrt{\pi_i / \|\boldsymbol{\pi}\|_1}$

as

$$\left[\frac{d_i - \mathbb{E}(d_i | \boldsymbol{\pi})}{\sqrt{\mathbb{E}(d_i | \boldsymbol{\pi})}} + \frac{\mathbb{E}(d_i | \boldsymbol{\pi}) - \pi_i \sqrt{\|\mathbf{d}\|_1}}{\sqrt{\mathbb{E}(d_i | \boldsymbol{\pi})}} \right] \frac{\|\boldsymbol{\pi}\|_1}{\sqrt{\|\mathbf{d}\|_1}} \sqrt{\frac{\mathbb{E}(d_i | \boldsymbol{\pi})}{\pi_i \|\boldsymbol{\pi}\|_1}}.$$

We must show that the first ratio above converges in distribution to a standard Normal, the second ratio converges in probability to zero, and the final multiplicative term to one; Slutsky's theorem then implies the result. To deduce the first result we use the Lindeberg–Feller central limit theorem as in Theorem 1. The second result follows from the delta method and Chebyshev's inequality. The third result follows by applying Chebyshev's inequality to $\|\mathbf{d}\|_1 / \|\boldsymbol{\pi}\|_1^2$ to deduce convergence in probability to 1, and then appealing to the continuous mapping theorem to obtain the same result for $\sqrt{\|\mathbf{d}\|_1} / \|\boldsymbol{\pi}\|_1$. $\square$

If we restrict $\gamma \in (0, 1/2)$ and $\theta_n = \omega(\sqrt{\log n} / n^{1/2-\gamma})$ as in Theorem 1, then all degrees will grow in n , and consequently this central limit theorem will hold for all elements of $\hat{\boldsymbol{\pi}} = \|\mathbf{d}\|_1^{-1/2} \mathbf{d}$.

2 A Hierarchical Model for Network Populations

2.1 From Conditional to Marginal Degree Distributions

An important shortcoming of the Poisson–Binomial model $\mathbb{P}(d_i = k | \boldsymbol{\pi})$ is that it cannot give rise to a heterogeneous *population* of networks, but instead is limited to random network realizations from a fixed and pre-specified vector $\boldsymbol{\pi}$ of normalized weights. The simplest construction giving rise to population heterogeneity will allow $\boldsymbol{\pi}$ itself to be random, resulting in a *hierarchical* generative model and a *marginal* degree distribution $\mathbb{P}(d_i = k) = \int \mathbb{P}(d_i = k | \boldsymbol{\pi}) f(\boldsymbol{\pi}) d\boldsymbol{\pi}$ for some probability distribution $f(\boldsymbol{\pi})$. If we further assume each π_i to be independent and identically distributed according to a probability distribution $f(\pi)$, then each d_i will be identically distributed as follows.

Proposition 1 (Hierarchical random graph properties). *Let $\pi_1, \dots, \pi_n$ be a random sample from some probability distribution $f(\pi)$ on $[0, 1]$, and let $\mathbf{d}$ be the degrees from an n -node simple graph whose edges are Bernoulli($\pi_i \pi_j$) trials. Then every element of $\mathbf{d}$ is identically distributed, taking values $k \in \{0, 1, \dots, n-1\}$ as follows:*

$$\mathbb{P}(d = k) = \binom{n-1}{k} \mathbb{E} \left\{ (\mathbb{E}(\pi) \pi)^k (1 - \mathbb{E}(\pi) \pi)^{n-1-k} \right\}, \quad (8)$$

$$\mathbb{E}(d) = (n-1) \mathbb{E}(\pi)^2, \quad (9)$$

$$\text{Var}(d) = (n-1) \mathbb{E}(d) \left[\text{Var}(\pi) + \frac{1 - \mathbb{E}(\pi^2)}{n-1} \right], \quad (10)$$

$$\text{Cov}(d_i, d_j) = \mathbb{E}(d) \left[\frac{3(n-2)}{n-1} \text{Var}(\pi) + \frac{1 - \mathbb{E}(\pi^2)}{n-1} \right], \quad i \neq j. \quad (11)$$

Proof. Observe that (2) is invariant to permutation of its parameters, and thus $\mathbb{E}_\pi [\mathbb{P}(d_i = k | \pi)] = \mathbb{E}_\pi \left[\binom{n-1}{k} \left(\prod_{j=1, j \neq i}^k \pi_i \pi_j \right) \left(\prod_{j=k+1, j \neq i}^{n-1} (1 - \pi_i \pi_j) \right) \right]$. For any i , the marginal expression in (8) then follows by evaluating scalar expectations with respect to each $\pi_{j, j \neq i}$. The moments of (9)–(11) follow by marginalizing their conditional counterparts in (4)–(6), using the law of total covariance in the latter two cases. $\square$

2.2 Reproducing Properties of Network Degrees

To understand the properties of observed network degrees under the hierarchical model described by Proposition 1, we must determine how our choice of $f(\pi)$ interacts with the sampling giving rise to $\mathbf{d}$, as characterized by the Poisson–Binomial distribution of (2) via (8). For well-behaved densities $f(\pi)$, the following theorem characterizes this interaction.

Theorem 3 (Reproducing property). *Assume $f(\pi)$ is bounded and everywhere nonzero on $[0, 1]$, with three bounded derivatives. Then*

$$n \mathbb{E}(\pi) \cdot \mathbb{P}(d=k) = f\left(\min\left[1, \frac{k}{n \mathbb{E}(\pi)}\right]\right) [1 - \tilde{F}(k; n, \mathbb{E}(\pi))] + \mathcal{E}(n),$$

where $\tilde{F}(k; n, \mathbb{E}(\pi))$ is the cumulative distribution function of a $\text{Binomial}(n, \mathbb{E}(\pi))$ random variable. The error term $\mathcal{E}(n)$ is

$$\mathcal{E}(n) = \begin{cases} \mathcal{O}\left(\frac{n-k}{n(n+1)}\right), & 0 < k \leq n \mathbb{E}(\pi); \\ \mathcal{O}\left(\frac{1}{n^{1/2-\epsilon}}\right) f(1) [1 - \tilde{F}(k; n, \mathbb{E}(\pi))], & n \mathbb{E}(\pi) < k < n. \end{cases}$$

If $f(\pi) = \text{Uniform}(0, 1)$, then $\mathcal{E}(n)$ is identically zero and

$$n \mathbb{E}(\pi) \cdot \mathbb{P}(d=k) = 1 - \tilde{F}(k; n, \mathbb{E}(\pi)). \quad (12)$$

Proof. We prove (12) and outline the general case for $f(\pi)$ satisfying the theorem hypotheses. Replace $\mathbb{E}(\pi)\pi$ by π in (8) to obtain

$$\mathbb{P}(d=k) = \frac{\binom{n-1}{k}}{\mathbb{E}(\pi)} \int_0^{\mathbb{E}(\pi)} \pi^k (1-\pi)^{n-k-1} f\left(\frac{\pi}{\mathbb{E}(\pi)}\right) d\pi; \quad (13)$$

this is the form of a mixed binomial [19], except for the appearance of $\mathbb{E}(\pi)$ as the upper limit of integration. If $f(\pi) = 1$ we immediately obtain (12), as in this case the integral of (13) evaluates to the product of the regularized incomplete Beta function $I_{\mathbb{E}(\pi)}(k+1, n-k) = 1 - \tilde{F}(k; n, \mathbb{E}(\pi))$ and the Beta function $\beta(k+1, n-k)$.

For the general case we proceed for each k via a Taylor series expansion of $f(\pi)$ about $\min[1, k/(n \mathbb{E}(\pi))]$, splitting the range of integration of (13) for each tail $k \leq n \mathbb{E}(\pi)$ according to $|k - n \mathbb{E}(\pi)| = \omega(\sqrt{n})$ or $|k - n \mathbb{E}(\pi)| = \mathcal{O}(\sqrt{n})$. For each Taylor term, the integrals over respective regions evaluate to differences of incomplete Beta functions. By choosing the regions as indicated, we

can use the boundedness of $f(\pi)$ and Hoeffding's inequality $\tilde{F}(k; n, \mathbb{E}(\pi)) \leq 0.5 \exp(-2(n \mathbb{E}(\pi) - k)^2/n)$ for $k \leq n \mathbb{E}(\pi)$ to show that most of these terms converge exponentially in n to 0 or a nonzero constant.

We are then left with the main term of (12) plus corresponding error terms, additive in the left tail (when $k \leq n \mathbb{E}(\pi)$) and multiplicative in the right (where $n \mathbb{E}(\pi) \mathbb{P}(d = k)$ is going rapidly to zero). Repeated application of this argument, combined with Lagrange's form of the Taylor remainder and the boundedness of the first three derivatives of $f(\pi)$, yields $\mathcal{E}(n)$ as stated. $\square$

Thus Theorem 3 fully characterizes degrees from independent Bernoulli($\pi_i \pi_j$) edges whenever the normalized weights $\pi_1, \dots, \pi_n$ comprise a random sample from some smooth and bounded density $f(\pi)$, nonzero on the entire unit interval $[0, 1]$. In particular, we see directly that any such $f(\pi)$ is *reproduced* in its entirety on the interval $0 \leq k \leq n \mathbb{E}(\pi)$.

This result also confirms that the marginal degree distribution $\mathbb{P}(d = k)$ results from the multiplicative interaction of $f(\pi)$ with the function $1 - \tilde{F}(k; n, \mathbb{E}(\pi))$, which is strictly decreasing in n and modulates $f(\min[1, k/(n \mathbb{E}(\pi))])$ over the entire range $0 \leq k \leq n - 1$ of d . As n grows large, the scaled probability mass $n \mathbb{E}(\pi) \mathbb{P}(d = k)$ is asymptotically equal to the product of these terms, the latter of which is the survival function of a Binomial($n, \mathbb{E}(\pi)$) random variable; i.e., the probability that such a random variable exceeds k . When $|k - n \mathbb{E}(\pi)| = \omega(\sqrt{n})$, this probability tends exponentially quickly to 1 or 0 by the arguments in the proof above, whereas in the transition region $|k - n \mathbb{E}(\pi)| = \mathcal{O}(\sqrt{n})$ we obtain a scaling between 0 and 1, attaining 1/2 as k approaches $n \mathbb{E}(\pi)$. The leftmost panels of Fig. 2 illustrate this behavior when $f(\pi) = \text{Uniform}(0, 1)$ in accordance with the exact expression of (12), with the remaining panels exemplifying the behavior of other smooth densities as indicated.

2.3 Networks from Scale-Free Distributions

An important example to which Theorem 3 does *not* directly apply is that of a scale-free or power-law density $f(\pi) \propto \pi^{-\beta}$, as this density is unbounded near zero. However, just as Theorem 1 demonstrated that fixing $\pi_i = i^{-\gamma}$ for $\gamma \in (0, 1/2)$ led asymptotically to the power law $\mathbb{P}(d = k) \propto k^{-\beta}$, where $\beta = 1 + 1/\gamma$, we show that such behavior is also achievable via a reproducing property for power-law densities.

Theorem 4 (Power law reproducing property). *Fix some $\beta \geq 0$, and let $\pi_1, \dots, \pi_n$ be drawn independently from $f(\pi) \propto \pi^{-\beta}$, restricted to the range $[l, u] \subseteq [0, 1]$, with $l > 0$ whenever $\beta \geq 1$. Then for all $k \in \{0, 1, \dots, n - 1\}$ such that $k > \beta - 1$,*

$$n \mathbb{E}(\pi) \cdot \mathbb{P}(d = k) = f\left(\frac{k+1}{(n+1) \mathbb{E}(\pi)}\right) \left[I_{u \mathbb{E}(\pi)}(k+1-\beta, n-k) - I_{l \mathbb{E}(\pi)}(k+1-\beta, n-k) \right] \left[1 + \mathcal{O}\left(\frac{(n-k)}{(n+1-\beta)(k+1-\beta)}\right) \right].$$




Figure 2: Simulated empirical degree distributions illustrating the reproducing property of Theorem 3, whereby $f(\pi)$ (top row) is reproduced for $\mathbb{P}(d = k)$ on the interval $0 \leq k \leq n \mathbb{E}(\pi)$ (bottom row; $n = 500$). Note the right tail behavior apparent in the bottom row, due to the multiplicative effect of $1 - \hat{F}(k; n, \mathbb{E}(\pi))$.

Proof. Let $c^{-1} = \int_l^u \pi^{-\beta} d\pi$, and note that when $\beta \geq 1$, we must have $l > 0$ to ensure integrability. We substitute $f(\pi/\mathbb{E}(\pi)) = c(\pi/\mathbb{E}(\pi))^{-\beta}$ in (13), adjusting the limits of integration to $l\mathbb{E}(\pi)$ and $u\mathbb{E}(\pi)$, respectively. For $k > \beta - 1$, we may then write the result as the difference of two incomplete Beta function integrals, yielding

$$\mathbb{P}(d = k) = \frac{c}{n \mathbb{E}(\pi)^{1-\beta}} \frac{\Gamma(n+1)}{\Gamma(n+1-\beta)} \frac{\Gamma(k+1-\beta)}{\Gamma(k+1)} \cdot [I_{u\mathbb{E}(\pi)}(k+1-\beta, n-k) - I_{l\mathbb{E}(\pi)}(k+1-\beta, n-k)]. \quad (14)$$

From the standard Taylor series for $\ln \Gamma(z)$, it is easily verified that $\Gamma(z)/\Gamma(z-\beta) = z^\beta [1 - (\beta/2)(1+\beta)(z-\beta)^{-1} + \mathcal{O}((z-\beta)^{-2})]$. Applying this to (14), first with $z = n+1$, and then with $z = k+1$ via the Taylor series of $\Gamma(z-\beta)/\Gamma(z)$, yields the stated result. $\square$

Theorem 4 shows that the scale-free density $f(\pi) \propto \pi^{-\beta}$ leads to a power-law decay in observed degree sequences across a range of k . For $k \leq \beta - 1$, it can be verified directly that the probability of observing $d = k$ decays exponentially in n . Similarly, when $k > u\mathbb{E}(\pi)n$, an exponential tail decay due to $I_{u\mathbb{E}(\pi)}(k+1-\beta, n-k)$ will rapidly take over. In fact, by analogy to the proof of Theorem 3, the difference $I_{u\mathbb{E}(\pi)}(k+1-\beta, n-k) - I_{l\mathbb{E}(\pi)}(k+1-\beta, n-k)$ goes




Figure 3: Left panel: Averaged degrees from power-law networks on $n = 1000$ nodes, generated from $f(\pi) \propto \pi^{-3}$ on $[1/3, 1]$, with $\mathbb{E}(\pi) = 1/2$. Rapid decays are visible near $k = 500/3 = l\mathbb{E}(\pi)n$ and $k = 500 = u\mathbb{E}(\pi)n$. Right panel: a log-log plot of the transition region near $\ln(u\mathbb{E}(\pi)n) \cong 6.2$, showing the exact and empirical distributions along with a Taylor expansion of $I_{u\mathbb{E}(\pi)}(k+1-\beta, n-k)$, corresponding to an exponential cutoff effect.

exponentially quickly to 0 or 1, with transition regions for $|l\mathbb{E}(\pi)n - k| = \mathcal{O}(\sqrt{n})$ and $|u\mathbb{E}(\pi)n - k| = \mathcal{O}(\sqrt{n})$.

Investigating this behavior in more detail reveals that the effects of Bernoulli sampling can explain power laws with exponential cutoff, which in practice often provide a better fit to data [20]. Power laws are typically identified from data by taking logarithms of empirical frequencies approximating $\mathbb{P}(d = k)$, and identifying a linear trend. Theorem 4 shows that $\ln \mathbb{P}(d = k)$ decomposes as a sum of contributions from $\ln[f((k+1)/[(n+1)\mathbb{E}(\pi)])]$ and $\ln[I_{u\mathbb{E}(\pi)}(k+1-\beta, n-k)]$. The latter term will serve to rapidly reduce probability mass as k exceeds $u\mathbb{E}(\pi)n$, and it is straightforward to show that a first-order Taylor expansion of this term corresponds to an exponential cutoff effect.

Figure 3 illustrates the consequence of this sampling-induced decay. The left panel shows $\mathbb{P}(d = k)$ and its empirical estimate for the case $\beta = 3$, with the right panel showing the transition region near $n\mathbb{E}(\pi)u$ on a logarithmic scale, along with a first-order Taylor expansion about the center of this region. This sounds a cautionary note to practitioners, in that effects ascribed to a model of degree sequence behavior may in fact be due purely to the effects of sampling.

3 Scaling and Sequences of Networks

3.1 Asymptotic Scaling Regimes

Theorems 3 and 4 confirm that marginal properties of degrees from independent Bernoulli($\pi_i\pi_j$) edges are determined by an interaction between a chosen distribution $f(\pi)$, and the Poisson–Binomial sampling distribution of (2). To this end, Fig. 4 shows four different distributions $f(\pi)$, each with mean $\mathbb{E}(\pi) = 1/2$,




Figure 4: Estimated values $\hat{p}_{ij} = d_i d_j / \|\mathbf{d}\|_1$ of $p_{ij} = \pi_i \pi_j$, sorted from largest to smallest degree to emphasize decay. Estimates are averaged over 10 network realizations for each indicated distribution $f(\pi)$, with $\mathbb{E}(\pi) = 1/2$ in each case.

in terms of estimates $\hat{p}_{ij} = d_i d_j / \|\mathbf{d}\|_1$, sorted from largest to smallest degree. The first corresponds to the scale-free example $f(\pi) \propto \pi^{-3}$ considered in Fig. 3 and Theorem 4; the second to the $\text{Uniform}(0, 1)$ example of Theorem 3; the third to a U-shaped distribution $\text{Beta}(1/2, 1/2)$; and the fourth an Erdős–Rényi graph with $p_{ij} = 1/2$. These distributions exemplify the wealth of structure that can be reproduced in this modeling regime. A key feature is that, despite their being standardized to have the same expectation, the resulting degrees all still retain distinctive structural properties.

This example motivates the question of precisely what ranges of limiting degree distributions are achievable by hierarchical simple random graph models. As long as elements of $\boldsymbol{\pi}$ remain order one, the combined effects of the reproducing property and Bernoulli sampling, as characterized by Theorems 3 and 4, will give rise to the behaviors above. Real networks exhibit substantial heterogeneity in their degree distributions, however, and so we must also understand modeling regimes in which these two effects are suitably balanced. This will enable us to construct degree distributions that scale in a controlled way with n , and exhibit greater heterogeneity than is possible when the order of $\boldsymbol{\pi}$

remains constant in n . For this reason, we introduce the notion of an *asymptotic scaling regime*, whose properties follow directly from (9)–(11) in Proposition 1.

Proposition 2 (Asymptotic scaling regime). *For π a random variable on $[0, 1]$ with fixed mean $\mathbb{E}(\pi)$ and variance $\text{Var}(\pi)$, we define an asymptotic scaling regime by the linear map $\pi^{(n)}(\pi)$, representing an n -dependent scaling and shifting of π as follows:*

$$\pi^{(n)}(\pi) = \theta_n \pi + \nu_n; \quad \theta_n = \frac{c_\theta}{n^{b_\theta}}, \quad \nu_n = \frac{c_\nu}{n^{b_\nu}}, \quad (15)$$

parameterized by $(b_\theta, c_\theta), (b_\nu, c_\nu)$, all strictly greater than zero.

Let $b_{\min} = \min(b_\nu, b_\theta)$. Degrees $d^{(n)}$ generated from $\pi^{(n)}(\pi)$ have mean and variance as follows, with $\mathcal{E}^{(n)} = \mathcal{O}(n^{-2b_{\min}})$:

$$\mathbb{E}(d^{(n)}) = n^{1-2b_\theta} (c_\theta \mathbb{E}(\pi) + c_\nu n^{b_\theta-b_\nu})^2 + \mathcal{E}^{(n)} \quad (16)$$

$$\begin{aligned} \text{Var}(d^{(n)}) &= n^{2(1-2b_\theta)} (c_\theta \mathbb{E}(\pi) + c_\nu n^{b_\theta-b_\nu})^2 \\ &\quad \cdot [c_\theta^2 \text{Var}(\pi) + n^{2b_\theta-1}] + n(\mathcal{E}^{(n)})^2. \end{aligned} \quad (17)$$

Note the dependence of $\mathbb{E}(d^{(n)})$ and $\text{Var}(d^{(n)})$ on n in (16) and (17), respectively. Beyond the case of fixed weights scaled by θ_n , as considered in Theorem 1, this n -dependence allows us to quantify the basic properties of $d^{(n)}$ in any asymptotic scaling regime, as we can shift and scale π independently via θ_n and ν_n . We observe directly from (16) and (17) that two regimes stand out as particularly important: the balanced asymptotic scaling regime in which $b_\theta = b_\nu$, whereupon the sampling variation in $d^{(n)}$ balances with the variability in π ; and the case of more limited heterogeneity in which $b_{\min} = 1/2$, so that both the mean and variance of each $d^{(n)}$ will converge to a finite limit in increasing n .

The expressions in (16) and (17) also highlight that a natural rescaling of $d^{(n)}$ is given by $1/n^{1-2b_{\min}}$. This is reminiscent of the deterministic construction of Theorem 1, for which an appropriate rescaling is $1/n^{1-2\gamma}$ for $\gamma < 1/2$. In Proposition 2 the overall magnitudes of the degrees can grow in a fashion commensurate to Theorem 1, but the degree variances will differ because π is random.

3.2 Networks with Controlled Growth of Expected Degrees

We start by exploring asymptotic scaling regimes in which $b_{\min}$, the slowest of the shift and scale decay rates, lies in $(0, 1/2)$. In this regime, each expected degree grows with network size n , but the expected *proportion* of nodes that a single node will connect to, compared to the $n-1$ possible, is decaying as $n^{-2b_{\min}}$. The total number of network edges will grow as $n^{2(1-b_{\min})}$, thus determining overall network density.

If $b_\nu \rightarrow \infty$ in this regime, so that the shift term ν_n goes to zero, we will achieve a reproducing property akin to Theorem 3. This can be shown by adapting a similar Taylor series argument to the magnitude of the derivatives of $f(c_\theta \pi / n^{b_\theta})$, with the error terms adjusted commensurately. As described above, the natural rescaling of each $d^{(n)}$ is then given by $1/n^{1-2b_\theta}$, so that the resultant random variables have order-one moments, yet inherit the properties of π .

3.3 Networks with Restricted Heterogeneity in Limiting Degrees

In certain networks it may be reasonable to expect that degree heterogeneity has saturated, meaning that (unlike in the case of controlled degree growth considered above) the distribution of degrees will not change appreciably as the network scales to larger sizes. This scenario implies a distribution of $d^{(n)}$ that is *independent* of n , and for which neither the mean nor the variance of each $d^{(n)}$ depends on n .

From (16) and (17), we see that this effect is achievable if $b_{\min} = 1/2$. A classic example is given by an Erdős–Rényi graph in the limiting Poisson regime: starting with an arbitrary $f(\pi)$ on $[0, 1]$, if we let $b_\theta \rightarrow \infty$ so that the effect of $f(\pi)$ is negligible, then for $b_\nu = 1/2$ we will obtain a Poisson degree distribution in the limit as $n \rightarrow \infty$. This is because $\pi^{(n)}$ will concentrate at $c_\nu/\sqrt{n}$, and hence each p_{ij} will scale as c_ν^2/n , so that the distribution of each $d^{(n)}$ approaches that of a Binomial($n - 1, c_\nu^2/n$) random variable.

Importantly, we may also achieve non-Poisson limits in this regime. Observe that if $b_{\min} = 1/2$, then the dispersion $\text{Var}(d^{(n)})/\mathbb{E}(d^{(n)})$ is given to leading order by $1 + c_\theta^2 \text{Var}(\pi)/n^{1-2b_\theta}$. This quantity converges to 1 when $b_\theta > 1/2$, but when $b_\theta = 1/2$, it tends to $1 + c_\theta^2 \text{Var}(\pi)$. Since the dispersion of a Poisson random variable is unity, we immediately see that when $b_\theta = 1/2$, the limiting distribution of $d^{(n)}$ cannot be Poisson.

The product $c_\theta^2 \text{Var}(\pi)$ is important in this non-Poisson setting, as it characterizes the limiting over-dispersion of $d^{(n)}$ relative to a Poisson variate, and will in turn govern limiting properties of the degrees. Note that c_θ^2 and $\text{Var}(\pi)$ cannot both be determined from a single network observation (though their product can); instead, to estimate them from data would require multiple realizations of the same network generating mechanism at different sample sizes n .

3.4 An Over-Dispersed Degree Limit

We conclude with a final example of degree heterogeneity that contrasts the Erdős–Rényi Poisson limit resulting from $b_\nu = 1/2$ and $b_\theta \rightarrow \infty$. There $\pi^{(n)}$ behaved like $c_\nu/\sqrt{n}$ as $n \rightarrow \infty$, and $f(\pi)$ had negligible effects. If we consider the *opposite* scenario of $b_\nu \rightarrow \infty$ and $b_\theta = 1/2$, and take the entropy-maximizing Uniform(0, 1) distribution, then the variation in $\pi^{(n)} = c_\theta \pi / \sqrt{n}$ will give rise to the following over-dispersed limit.

Proposition 3 (An over-dispersed degree limit). *Let $f(\pi) = 1$ on $[0, 1]$, and define $\pi^{(n)}(\pi)$ via the asymptotic scaling regime of (15), with $b_\nu \rightarrow \infty$ and $b_\theta = 1/2$. Then as $n \rightarrow \infty$, each degree $d^{(n)}$ converges to a random variable $d^{(\infty)}$ such that for $\zeta = c_\theta^2/2$, we have $\mathbb{E}(d^{(\infty)}) = \zeta/2$, $\text{Var}(d^{(\infty)}) = \zeta(\zeta + 6)/12$, and*

$$\mathbb{P}(d^{(\infty)} = k) = \frac{1}{\zeta} \left(1 - \sum_{j=0}^k \frac{e^{-\zeta} \zeta^j}{j!} \right), \quad k = 0, 1, \dots$$

Proof. We start from (8), which gives a general expression for the marginal probability $\mathbb{P}(d^{(n)} = k)$. Two different approaches must be adopted: one for $k = o(n)$, and the other for $k \propto n$. In the former scenario, we evaluate the limit of the expression $(1 - \mathbb{E}(\pi)\pi)^{n-k-1}$ from (8) using $\lim_{n \rightarrow \infty} (1 - x/n)^n = e^{-x}$, and evaluate the resulting integral directly. For $k \propto n$, the probability of obtaining k decays exponentially in n . If we bound (8) in this regime using Stirling's inequality, the triangle inequality can then be used to obtain convergence to the stated $\mathbb{P}(d^{(\infty)} = k)$. The mean and variance of this limiting distribution are then calculated directly. $\square$

Observe from the above that $\pi^{(n)}$ is a uniform variate whose range $[0, c_\theta/\sqrt{n}]$ is shrinking toward zero at rate $1/\sqrt{n}$, without any significant shift parameter ν_n to offset the reduction in its mean. Thus, as we expect, Proposition 3 verifies that $\mathbb{P}(d^{(\infty)} = k)$ decays monotonically as k increases from zero. Yet $d^{(\infty)}$ is over-dispersed relative to a Poisson variate, as $\text{Var}(d^{(\infty)})/\mathbb{E}(d^{(\infty)}) = 1 + \mathbb{E}(d^{(\infty)})/3$. Since the dispersion of $d^{(\infty)}$ depends linearly on its mean $\mathbb{E}(d^{(\infty)}) = c_\theta^2/4$, it follows that for increasingly larger values of c_θ , we can expect to see increasingly variable realizations of $d^{(\infty)}$ relative to a Poisson limiting regime. In this same manner, other asymptotic scaling regimes can yield different limiting behaviors.

4 Discussion

In this article we have introduced a hierarchical statistical model of simple random graphs, and used it to give a precise answer to the question, “What is a degree distribution?” We have developed exact statistical results, limit theorems, and large-sample approximations that govern the behavior of networks based on our construction. The advantage of using a statistical model to do so is that it links networks of varying sample sizes to a single range of parameters, and indicates under what assumptions these parameters can be estimated.

To study a fully generative modeling framework, we placed a stochastic model on normalized weights giving rise to a network adjacency matrix. This enabled us to understand the notions of distributions of degrees *between* networks, as well as *within* the context of a single network. We determined a reproducing property associated with these normalized weights, and explicitly quantified effects for finite but large n —including explaining the exponential cutoff behavior often noted in this setting as an effect of Bernoulli sampling. Finally,

to understand explicitly the effects of sampling variability, we explored various asymptotic scaling regimes. These permitted us to scale order-one weights toward zero as $n \rightarrow \infty$, thereby allowing for the control of limiting degree distributions across network sizes.

References

- [1] P. J. Bickel and A. Chen. A nonparametric view of network models and Newman–Girvan and other modularities. *Proc. Natl. Acad. Sci. USA*, 106:21068–21073, 2009.
- [2] S. E. Fienberg and A. Rinaldo. Maximum likelihood estimation in log-linear models. *Ann. Statist.*, 40:996–1023, 2012.
- [3] M. Girvan and M. E. J. Newman. Community structure in social and biological networks. *Proc. Natl. Acad. Sci. USA*, 99:7821–7826, 2002.
- [4] D. Lazer et al. Life in the network: the coming age of computational social science. *Science*, 323:721–723, 2009.
- [5] A. L. Barabási and R. Albert. Emergence of scaling in random networks. *Science*, 286:509–512, 1999.
- [6] M. E. J. Newman, D. J. Watts, and S. H. Strogatz. Random graph models of social networks. *Proc. Natl. Acad. Sci. USA*, 99:2566–2572, 2002.
- [7] T. Britton, M. Deijfen, and A. Martin-Löf. Generating simple random graphs with prescribed degree distribution. *J. Statist. Phys.*, 124:1377–1397, 2006.
- [8] B. Bollobás, S. Janson, and O. Riordan. The phase transition in inhomogeneous random graphs. *Random Struct. Alg.*, 31:3–122, 2007.
- [9] M. Molloy and B. Reed. A critical point for random graphs with a given degree sequence. *Rand. Struct. Alg.*, 6:161–180, 1995.
- [10] R. Durrett. *Random Graph Dynamics*. Cambridge U. Pr., 2007.
- [11] F. Chung and L. Lu. The average distances in random graphs with given expected degrees. *Proc. Natl. Acad. Sci. USA*, 99:15879–15882, 2002.
- [12] P. O. Perry and P. J. Wolfe. Null models for network data. arXiv:1201.5871, 2012.
- [13] S. Janson. Asymptotic equivalence and contiguity of some random graphs. *Random Struct. Alg.*, 36:26–45, 2010.
- [14] R. van der Hofstad. Critical behavior in inhomogeneous random graphs. *Random Struct. Alg.*, 2012. In press.

- [15] S. Chatterjee, P. Diaconis, and A. Sly. Random graphs with a given degree sequence. *Ann. Appl. Probabil.*, 21:1400–1435, 2011.
- [16] M. E. J. Newman. Modularity and community structure in networks. *Proc. Natl. Acad. Sci. USA*, 103:8577–8582, 2006.
- [17] F. Chung, L. Lu, and V. Vu. Spectra of random graphs with given expected degrees. *Proc. Natl. Acad. Sci. USA*, 100:6313–6318, 2003.
- [18] A. D. Barbour, L. Holst, and S. Janson. *Poisson Approximation*. Oxford U. Pr., 1992.
- [19] A. Hald. The mixed binomial distribution and the posterior distribution of p for a continuous prior distribution. *J. Roy. Stat. Soc. B*, pages 359–367, 1968.
- [20] M. E. J. Newman. The structure of scientific collaboration networks. *Proc. Natl. Acad. Sci. USA*, 98:404–409, 2001.